

# Songbird ReMix **BOP v1 Extras**

for Songbird Remix (available through Hivewire3D [www. Hivewire3D.com](http://www.Hivewire3D.com))  
Characters & Textures by Ken Gilliland ([www.empken.com](http://www.empken.com))

For all Songbird ReMix information and free downloads visit: [www.songbirdremix.com](http://www.songbirdremix.com)

**LICENSE AGREEMENT:** By downloading these files you agree not to sell them, redistribute them in any compilations or collections or any derivatives thereof, to anyone, without the express written consent of the author, Ken Gilliland. Using 10% or more of the texture maps for the purposes of creating a new product violates this license agreement.

## **Required:**

- Songbird ReMix Birds of Prey Volume 1: Kestrels, Hobbys & Falcons

## **Installation for Poser 8+:**

- Copy the "Runtime" folder to your Poser folder.

## **Installation for Native DAZ|Studio v4.5+:**

- Copy the "Content" folder to your DAZ|Studio Content folder.

## **Includes:**

- *Peregrine Falcons*
  - *F. p. brookei*, described by Sharpe in 1873. It is also known as the Mediterranean peregrine falcon or the Maltese falcon. It includes caucasicus and most specimens of the proposed race punicus, though others may be pelegrinoides, barbary falcons, or perhaps the rare hybrids between these two which might occur around Algeria. They occur from the Iberian Peninsula around the Mediterranean, except in arid regions, to the Caucasus. They are non-migratory. It is smaller than the nominate subspecies, and the underside usually has rusty hue. Males weigh around 445 grams (0.981 lb), while females weigh up to 920 grams (2.03 lb).
  - *F. p. ernesti*, described by Sharpe in 1894. It is found from Indonesia to Philippines and south to Papua New Guinea and the nearby Bismarck Archipelago. Its geographical separation from *nesiotes* requires confirmation. It is non-migratory. It differs from the nominate subspecies in the very dark, dense barring on its underside and its black ear coverts.
  - *F. p. macropus*, described by Swainson in 1837. It is the Australian peregrine falcon. It is found in Australia in all regions except the southwest. It is non-migratory. It is similar to *brookei* in appearance, but is slightly smaller and the ear region is entirely black. The feet are proportionally large.
- *Seychelles Kestrel (Immature)*